



WHAT IS CLAIMED IS:

1. A compound having a structural formula I,




or a pharmaceutically acceptable salt, solvate, hydrate or stereoisomer thereof, wherein:
R¹ and R² are each independently: methyl or ethyl.

2. The compound of Claim 1, wherein the compound having a structural formula II,




or a pharmaceutically acceptable salt, solvate or hydrate thereof, wherein: R¹ and R² are
each independently: methyl or ethyl.

3. The compound of Claim 2, wherein the compound is (2S)-3-(4-{[2-(4-methoxy-phenyl)-ethylcarbamoyl]-methoxy}-phenyl)-2-methoxy-propionic acid
having a structural formula III,




or a pharmaceutically acceptable salt, solvate or hydrate thereof.

4. The compound of Claim 1, wherein the compound is 3-(4-{[2-(4-ethoxy-phenyl)-ethylcarbamoyl]-methoxy}-phenyl)-2-methoxy-propionic acid having a structural formula IV,



or a pharmaceutically acceptable salt, solvate or hydrate thereof.

5. The compound of Claim 4, wherein the compound is (*S*)-3-(4-{[2-(4-ethoxy-phenyl)-ethylcarbamoyl]-methoxy}-phenyl)-2-methoxy-propionic acid having a structural formula V,



or a pharmaceutically acceptable salt, solvate or hydrate thereof.

15

6. A pharmaceutical composition comprising a pharmaceutically acceptable carrier and a compound of Claims 1-5 or a pharmaceutically acceptable salt, solvate or hydrate thereof.

20

7. A pharmaceutical composition comprising:

(1) a compound of Claims 1-5, or a pharmaceutically acceptable salt, solvate, hydrate or stereoisomer thereof;

(2) a second therapeutic agent selected from the group consisting of: insulin sensitizers, sulfonylureas, biguanides, meglitinides, thiazolidinediones, α -

glucosidase inhibitors, insulin secretagogues, insulin, antihyperlipidemic agents, plasma HDL-raising agents, HMG-CoA reductase inhibitors, statins, acyl CoA:cholesterol acyltransferase inhibitors, antiobesity compounds, antihypercholesterolemic agents, fibrates, vitamins and aspirin; and

5 (3) optionally a pharmaceutically acceptable carrier.

8. A method of modulating a peroxisome proliferator activated receptor (PPAR) comprising the step of contacting the receptor with a compound of Claims 1-5, or a pharmaceutically acceptable salt, solvate or hydrate thereof.

10

9. The method of Claim 8, wherein the PPAR is an alpha (α)-receptor.

10. The method of Claim 8, wherein the PPAR is a gamma (γ)-receptor.

15

11. The method of Claim 8, wherein the PPAR is a alpha/gamma (α/γ)-receptor.

12. A method for treating a PPAR γ mediated disease or condition in a mammal comprising the step of administering an effective amount of a compound of Claims 1-5.

20

13. A method for treating a PPAR α mediated disease or condition in a mammal comprising the step of administering an effective amount of a compound of Claims 1-5.

25

14. A method for treating a PPAR α/γ mediated disease or condition in a mammal comprising the step of administering an effective amount of a compound of Claims 1-5.

30

15. A method for treating disease or condition mediated by a PPAR γ partial agonist in a mammal comprising the step of administering an effective amount of a compound of Claims 1-5.
- 5 16. A method for lowering blood-glucose in a mammal comprising the step of administering an effective amount of a compound of Claims 1-5.
- 10 17. A method of treating disease or condition in a mammal selected from the group consisting of hyperglycemia, dyslipidemia, Type II diabetes, Type I diabetes, hypertriglyceridemia, syndrome X, insulin resistance, heart failure, diabetic dyslipidemia, hyperlipidemia, hypercholesteremia, hypertension, obesity, anorexia bulimia, anorexia nervosa, cardiovascular disease and other diseases where insulin resistance is a component, comprising the step of administering an effective amount of a compound of Claims 1-5.
- 15 18. A method of treating diabetes mellitus in a mammal comprising the step of administering to a mammal a therapeutically effective amount of a compound of Claims 1-5.
- 20 19. A method of treating cardiovascular disease in a mammal comprising the step of administering to a mammal a therapeutically effective amount of a compound of Claims 1-5, or a pharmaceutically acceptable salt, solvate or hydrate thereof.
- 25 20. A method of treating syndrome X in a mammal, comprising the step of administering to the mammal a therapeutically effective amount of a compound of Claims 1-5, or a pharmaceutically acceptable salt, solvate or hydrate thereof.
- 30 21. A method of treating a disease or condition in a mammal selected from the group consisting of hyperglycemia, dyslipidemia, Type II diabetes, Type I diabetes, hypertriglyceridemia, syndrome X, insulin resistance, heart failure, diabetic dyslipidemia, hyperlipidemia, hypercholesteremia, hypertension, obesity, anorexia

- bulimia, anorexia nervosa, cardiovascular disease and other diseases where insulin resistance is a component, comprising the step of administering an effective amount of a compound of Claims 1-5; and an effective amount of second therapeutic agent selected from the group consisting of: insulin sensitizers, sulfonylureas, biguanides, meglitinides,
- 5 thiazolidinediones, α -glucosidase inhibitors, insulin secretagogues, insulin, antihyperlipidemic agents, plasma HDL-raising agents, HMG-CoA reductase inhibitors, statins, acyl CoA:cholesterol acyltransferase inhibitors, antiobesity compounds, antihypercholesterolemic agents, fibrates, vitamins and aspirin.
- 10 22. Use of a compound of Claims 1-5, or a pharmaceutically acceptable salt, solvate or hydrate thereof, for the manufacture of a medicament for the treatment of a condition modulated by a PPAR.
23. Use of a compound of Claims 1-5, or a pharmaceutically
- 15 acceptable salt, solvate or hydrate thereof, for the manufacture of a medicament for the treatment of diabetes.